

Begonia boliviensis

Santa Cruz® Sunset

Culture guide

Uses:

Annual – Ideal for hanging baskets, containers and garden landscape

Exposure:

Sun - Shade

Garden height:

12" / 30 cm

Crop time:

14-16 weeks

Sowing method:

1 seed per pellet. 288 tray recommended

Germination:

Stage I: 7-14 days at 70-73 °F (21-23 °C) and relative humidity of 95 %. Maintain high moisture. Light germinator, do not cover pellets. Additional light (50 W/m²) accelerates germination and improves quality of the seedlings

Stage II: 7-14 days at 68-70 °F (20-21 °C).

Continue long day lighting (14 h) and avoid saturated or dry media, lower humidity levels.

Water temperature should be 64 °F (18 °C)

Stage III-IV: 28-42 days at 64-68 °F (18-20 °C)

Temperature:

Grow at 60-64 °F (16-18 °C). 10 days before selling temperature can be decreased to 60 °F (16 °C). Temperatures below 57 °F (14 °C) will result in tuber formation and crop delay.

Fertilization:

Moderate fertilization levels are required. Fertilize weekly with 150-200 ppm

nitrogen, using a complete and potassium balanced fertilizer (N:K₂O-ratio: 1:1,5). Keep low ammonium levels, otherwise the roots become damaged. At high nitrogen levels the foliage can become very big. Avoid pH above 6.5, as high pH causes iron deficiency. Apply chelated iron, if chlorosis becomes a problem. To prevent magnesium deficiency apply magnesium sulphate (0,025 %) 1-2 times. Additional foliage fertilization with potassium supports compact plant

growth and provides a dark green foliage color.

Avoid high soluble salts in the soil.

Ref. : 8 May 2013

www.benary.com

