

GrowerFacts

Germination Approximate seed count: 18,425-21,000 S./oz. (650-750 S./g)

Media Use a well-drained, disease-free seedling medium with a pH of 5.8 to 6.2. Maintain EC of about 0.75 to 0.80 mS/cm (1:2 extraction).

Sowing Can be produced in a 288 cell tray or larger size plug tray at 1 seed per cell. Covering the seed is optional. A light cover can help maintain high relative humidity around the seed and improve germination.

Stage 1 Germination in approximately 3 to 6 days. Move plug trays out of germination chamber as soon as radicles emerge to avoid hypocotyl stretch. Soil temperature: 65 to 72°F (18 to 22°C) Note: Germinates readily at temperatures of 62 to 77° F (16 to 25°C). Warmer temperatures result in faster germination.

Light: Light is optional.

Moisture: Keep soil wet (level 4) during stage 1. Humidity: Maintain 95% or greater relative humidity (RH) until radicle emergence.

Plug Production

Stage 2

Soil temperatures: 65 to 72°F (18 to 22°C) Light: Up to 2,500 f.c. (26,900 Lux)

Moisture: Maintain high moisture (level 4).

Fertilizer: Apply fertilizer at rate 1 (less than 100 ppm N/less than 0.7 mS/cm EC) with a nitrate-form fertilizer with low phosphorus.

Stage 3

Soil temperatures: 65 to 72°F (18 to 22°C)

Light: Up to 2,500 f.c. (26,900 Lux)

Moisture: Soil moisture can be reduced to level 3. Keep wet-dry moisture cycle between levels 3 and 4. Avoid seedling wilting.

Fertilizer: Increase fertilizer to rate 2 (100 to 175 ppm N/ 0.7 to 1.2 mS/cm EC). Maintain medium pH 5.8 to 6.2 and EC between 1.0 and 1.5 mS/cm (1:2 extraction).

Stage 4

Soil temperatures: 65 to 72°F (18 to 22°C)

Light: Up to 5,000 f.c. (54,000 Lux).

Moisture: Moisture level can be reduced to medium to medium dry (level 3 to 2). Do not allow the seedlings to wilt.

Fertilizer: Same as stage 3

Growth regulators: B-Nine/Alar (daminozide) is effective for Glitz plug height control. Two applications of B-Nine/Alar 2500 ppm (3 g/l, 85% formulation or 4 g/l, 64% formulation) spray starting at true leaf initial stage (early stage 3), followed by a second application 7 days later, works well for plug height control. Adding sticker to the solution may cause leaf damage. Alternatively, a Bonzi (paclobutrazol) 2.5 ppm (0.6ml/l, 0,4% formulation) sprench or Bonzi 0.25 to 0.5 ppm (0.06 to 0.12 ml/l, 0,4% formulation) drench at radicle emergence stage has been shown effective to control hypocotyl stretch.

Pinching: Not recommended.

Note: Do not grow plugs at temperatures below 62°F (16°C) as it could cause leaves to yellow and drop.

Transplanting: Euphorbia Glitz plugs can be transplanted deeply to the level of the first node for stretched plugs.



EUPHORBIA *graminea* GLITZ

60-7439

Container Size 4 to 5-in (10 to 13 cm), quart pot (12 cm): 1 plug per pot

6-in (15 cm) pot: 2 plugs per pot

10-in (25 cm) HB: 3 plugs per pot

Media Use a well-drained, disease-free growing medium with a pH of 5.8 to 6.2.

Temperature nights: 65 to 68°F (18 to 20°C) **Days:** 65 to 77°F (18 to 25°C) Avoid temperatures below 62°F (16°C) as cooler temperatures cause foliage yellowing.

Light Keep light as high as possible while maintaining recommended temperature.

Media Moisture The medium should be allowed to dry between watering. Do not allow the plants to wilt.

Fertilizer Starting a week after transplant, apply fertilizer at rate 3 (175 to 225 ppm N/1.2 to 1.5 mS/cm) once a week from nitrate-form fertilizer with low phosphorus. Maintain the media EC at 1.5 to 2.0 mS/cm at pH at 5.8 to 6.2.

Pinching Pinching is not recommended as it can cause poor habit development and increase crop time.

Plant Growth Regulators Applications of B-Nine/Alar 2,500-5,000 ppm (3 to 6 g/l, 85% formulation or 4 to 8 g/l, 64% formulation) spray have been shown to be effective. Start the first application one week after transplant. Repeat as needed.

Note: It is the responsibility of the applicator to read and follow all current label directions for the specific chemical being used and to use the PGR in accordance with all laws and regulations.

Crop Scheduling

Germination: 3 to 6 days

Sow to Transplant (288 cell): 4 to 5 weeks

Transplant to finish:

Container Size: 4 to 5-in. (10 to 13-cm), quart pot

288-cell plugs per pot: 1

Weeks from transplant: 5 to 8

Total weeks*: 9 to 13

Container Size: 6-in. (15-cm) pot to 1 gallon

288-cell plugs per pot: 2

Weeks from transplant: 5 to 8

Total weeks*: 9 to 13

Container Size: 10-in. (25-cm) basket

288-cell plugs per pot: 3

Weeks from transplant: 6 to 8

Total weeks*: 10 to 13

Common Problems

Insect: Thrips, aphids, and spider mites.

Disease: No serious problems.

Landscape and garden performance

- Not recommended to transplant Glitz Euphorbia outdoors until temperatures are above 62°F (16°C).
- Works well solo or in mixed baskets, containers and patio planters.
- Garden location: Full sun to partial sun
- Garden spacing: 8 to 12 in. (20 to 30 cm)
- Finished height: 10 to 14 in. (25 to 35 cm)
- Finished spread: 12 to 18 in. (30 to 45 cm)

Ref: June 16, 2014

http://www.panamseed.com/plantlist.aspx?dispmode=grower_newvarieties



NORSECO