

Impatiens (Interspecific) Solarscape™ F₁ Series

Annuals Culture (revised 06/08/22)

(*Impatiens hybrida*)

Impatiens landscapers will love

NEW Solarscape seed-propagated interspecific impatiens provides unbeatable colour in full-sun garden beds and containers. Durable Solarscape performs just like vegetative varieties with the flexibility of seed. Landscape growers will appreciate this option for non-branded needs in-ground that grows strong in the sun!

Plug crop time: (288) 4 to 5 weeks, (128) 5 to 6 weeks

Transplant to finish: 7 to 10 weeks

Available in North America only.

- Great low-maintenance plant provides high-impact colour in full-sun spaces, whether in the ground or in containers.
- The first interspecific impatiens series from seed offers growers the flexibility of an input that is available all year round, making scheduling a breeze.
- Seed-propagated Solarscape provides the same garden performance as vegetative interspecific impatiens without the royalties! Perfect for opening-price-point, non-branded programs at retail.
- Offers landscape growers a non-branded, cost-efficient alternative.
- Blooms have tropical appeal, with a satiny texture in vibrant, glowing hues.

General Information

Exposure	Bloom Season	Height	Spread	Spacing
Partial Sun, Sun	Spring, Late Spring, Summer, Late Summer	9-11 in. (23-28 cm)	15-20 in. (38-51 cm)	12-14 in. (30-36 cm)

Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days from 50% to maximum germination	Initial Media pH/EC (1:2)	Cover Seed
RAW	288	1	4-5	4-5	5.8-6.2 pH 0.05-0.07 mmhos/cm	Light cover
	128	1	5-6			

Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 4-5	Level 4	Level 3-4	Level 2-4
Temperature	72-77°F (22-25°C)	72-77°F (22-25°C)	65-72°F (18-22°C)	65-72°F (18-22°C)
Light	Light	450-700 f.c. (4,800-7,500 Lux)	450-700 f.c. (4,800-7,500 Lux)	450-700 f.c. (4,800-7,500 Lux)
Fertilizer		Less than 100 ppm N (Less than 0.7 EC)	Less than 100 ppm N (Less than 0.7 EC)	Less than 100 ppm N (Less than 0.7 EC)
PGR		daminozide/2,500-5,000 ppm/Spray paclobutrazol/1 ppm/Spray	daminozide/2,500-5,000 ppm/Spray	

Propagation Key Tips

Moisture levels surrounding the seed are very important in stage 1 and 2. Light is required for optimal germination, so is high relative humidity. A light cover of vermiculite may be beneficial to raise humidity around the seed. White Shimmer may be 2 days slower. Avoid temperatures lower than 62°F/17°C. Temperatures at or above 85°F/29°C result in slower germination speed and lower germination results. In stage 3, repeat daminozide 2,500-5,000 ppm spray if necessary.

Growing on to Finish

Growing on Temperature	Target Media pH/EC (1:2)	Fertilizer	Daylength
(day) 68-75°F (20-24°C) (night) 60-68°F (16-20°C)	5.8-6.2 pH 0.5-0.75 mmhos/cm	Less than 100 ppm N (Less than 0.7 EC)	Day Neutral

Daylength Notation

Light Recommendations: 4,000-8,000 foot candles. Recommended DLI of 10 or greater moles. In areas of low DLI, less than 10 moles, supplemental lighting is beneficial to increase quantity and speed of flowering. In areas of low relative humidity, less than 30%, shading is recommended.

Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
306 Pack/1801	1 (ppp)	7-8 (weeks)	Spring	-
4"/4.5"/Quart/10 cm	1 (ppp)	7-8 (weeks)	Spring	-
5"/6"/1 Gallon/15 cm	1 (ppp)	8-9 (weeks)	Spring	-
8"/2 Gallon/20 cm	1-3 (ppp)	9-10 (weeks)	Spring	-

Fertilizer Notation

Solarscapes are low-to-moderate feeders, therefore low rates of the nitrate form of nitrogen fertilizer are recommended: 75-100 ppm N (EC target of 0.5 to 0.75).

Chemical Sensitivity

This product is very responsive to PGRs. Drenching is not recommended.

Common Problems

Insects: Thrips, Aphids, Fungus Gnats, Mites. Diseases: Pythium, Rhizoctonia, Phytophthora, Botrytis, Tomato Spotted Wilt Virus, Impatiens Necrotic Spot Virus, Powdery Mildew, Myrothecium

Finishing Key Tips

Temperature Recommendations: Minimum: 60°F/16°C Maximum: 85°F/29°C Lowering ADT by 2°F/1°C will delay flowering by 2 to 4 days, with an even longer delay at cooler ADTs. Moisture Recommendations: Plants are well controlled with water and fertilization management; growing dry with less to no PGRs results in plants that are faster to establish in the garden. PGR Recommendations: 10 to 14 days after transplant to finish container, repeating as needed. Avoid over application of PGRs, as this may result in plant stunting or slow growing-on. If growing in dry conditions, the lowest rates of PGRs are recommended. Solarscape Magenta Bliss, Orange Burst and White Shimmer: paclobutrazol 0.25 ppm drench OR paclobutrazol 5-10 ppm spray Solarscape Neon Purple (most compact hybrid): paclobutrazol 0.10 ppm drench OR 5 ppm spray

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

Variety Pictures



Magenta Bliss



Neon Purple



Orange Burst



White Shimmer

PanAmericanSeed™

PanAmerican Seed Co.
622 Town Road, West Chicago, Illinois, USA, 60185-2698
630 231-1400 Fax: 630 231-3609 PanAmSeed.com

™ denotes a trademark of and © denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.
©2022 Ball Horticultural Company