

## Dianthus (Interspecific) Jolt™ F<sub>1</sub> Series

(*Dianthus barbatus interspecific*)

Annuals Culture (revised 01/27/22)

**Plug crop time:** 4 to 5 weeks

**Transplant to finish:** Late Spring through Autumn finish, 11 to 13 weeks; Winter finish, 14 to 18 weeks

- *Shocking Summer Colour* from the most heat-tolerant Interspecific Dianthus!
- Electrifying colours and all-season performance make Jolt stand out in landscapes and premium containers.
- Highly durable plants withstand hot conditions with few disease issues and continue to flower without setting seed.
- Grower-friendly, well-branched plants are easy to produce with less to no PGRs.
- Perfect for Summer premium grower choice quarts and gallons to command a higher price point.
- Strong dark green foliage.
- Performs as a semi-hardy annual in Zone 7 (0°F/-18°C minimum temperature) and milder.
- *All-America Selections National Winner* Jolt Pink continues to stir up consumer excitement with extra publicity and promotion.

2019 Louisiana State University Super Plant

### General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	Spring, Late Spring, Summer, Late Summer, Autumn	16-20 in. (41-51 cm)	12-14 in. (30-36 cm)	10-14 in. (25-36 cm)

### Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days from 50% to maximum germination	Initial Media pH/EC (1:2)	Cover Seed
PEL	288	1	5-6	3-4	5.8-6.2 pH 0.75 mmhos/cm	Yes

### Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
<b>Moisture</b>	Level 4	Level 3-4	Level 2-3	Level 2-3
<b>Temperature</b>	64-68°F (18-20°C)	65-70°F (18-21°C)	60-65°F (16-18°C)	55-60°F (13-16°C)
<b>Light</b>	Optional	5-8 mol·m <sup>-2</sup> ·d <sup>-1</sup>	8-10 mol·m <sup>-2</sup> ·d <sup>-1</sup>	10 mol·m <sup>-2</sup> ·d <sup>-1</sup>
<b>Fertilizer</b>		Less than 100 ppm N (Less than 0.7 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	175 to 225 ppm N (1.2 to 1.5 EC)
<b>PGR</b>		paclobutrazol/5 ppm/Spray	paclobutrazol/5 ppm/Spray	

### Propagation Key Tips

A small percentage (3-5%) of flowering off-types can be observed with Jolt dianthus at 4 to 5 weeks from sowing. These plants should be removed/discarded at transplant. Jolt Purple is more sensitive to paclobutrazol than others and the rate should be reduced to half of others.

### Growing on to Finish

Growing on Temperature	Target Media pH/EC (1:2)	Fertilizer	Daylength
(day) 65-75°F (18-24°C) (night) 60°F (16°C)	5.8-6.2 pH 1.5-2.0 mmhos/cm	175 to 225 ppm N (1.2 to 1.5 EC)	Facultative Long Day

### Daylength Notation

Will flower under all photoperiods but ~1 week earlier at 15 hours (LD) than 10 hours (SD). At a given photoperiod, plants flower up to 1.5 weeks faster under high daily light integral than low.

## Crop Scheduling

---

Container Size	Plugs/Pot	Crop Time	Season	PGR
4"/4.5"/Quart/10 cm	1 (ppp)	14-18 (weeks)	Spring	paclobutrazol 20 ppm Spray
5"/6"/1 Gallon/15 cm	1-3 (ppp)	14-18 (weeks)	Spring	paclobutrazol 20 ppm Spray
4"/4.5"/Quart/10 cm	1 (ppp)	11-13 (weeks)	Summer	paclobutrazol 20 ppm Spray
5"/6"/1 Gallon/15 cm	1-3 (ppp)	11-13 (weeks)	Summer	paclobutrazol 20 ppm Spray

### Fertilizer Notation

Starting a week after transplant, apply fertilizer at rate 3 (175 to 225 ppm N/1.2 to 1.5 mS/cm) once a week from a nitrate-based fertilizer with low phosphorus.

### Common Problems

Insects: Thrips, Mites

### Finishing Key Tips

Temperature and light intensity have greater impact on flowering, especially during Winter and early Spring. Jolt will benefit from being grown under high light levels and is a facultative/quantitative long-day plant. It can flower under different daylengths, but will take slightly longer to flower under short days than long days. A small percentage (up to 3%) of early off-types can be observed with Jolt dianthus at 4 to 5 weeks from sowing. These plants should be removed/discarded at transplant. Jolt Purple is more sensitive to paclobutrazol than others and the rate should be reduced to half of others.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

## Variety Pictures

---



Cherry



Pink



Pink Magic



Purple

PanAmericanSeed™

PanAmerican Seed Co.  
622 Town Road, West Chicago, Illinois, USA, 60185-2698  
630 231-1400 Fax: 630 231-3609 PanAmSeed.com

™ denotes a trademark of and © denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.  
©2022 Ball Horticultural Company