

# *Kniphofia uvaria*

## **FLAMENCO**

Red Hot Poker, Torchlily

### **Culture guide**

**Uses:**

Evergreen perennial for border and bedding, cut flowers

**Exposure:**

Sun

**Garden height:**

31" / 80 cm

**Crop time:**

26-28 weeks

**Sow time:**

January-May for flowering in pots from July onwards, June-August for flowering in pots the following year

**Sowing method:**

2-3 seeds per plug, can be sown directly in final pots (4-6 seeds)

**Germination:**

Germinates in 14-20 days at 65-72 °F (18-22 °C).

Light is required for germination. During Stage II, maintain temperatures of 65-68 °F (18-20 °C).

During Stages II and IV, gradually reduce temperature to 58-62 °F (14-16 °C).

**Growing On:**

Transplant plugs after 8 weeks. Grow on at of 60 °F

(15 °C) in a well-drained, humus, loamy medium.

Vernalization is not required for flowering

**Media:**

Use a well-drained, growing perennial substrate with 0-15 % clay, 0-15 % parts (e.g. bark, wood fibres), 1,5-2 kg/m<sup>3</sup> complete balanced fertilizer, 2-3 kg/m<sup>3</sup> slow release fertilizer (3-9 months), iron-chelate, micronutrients, pH: 5.5-6.2. Field: humus sandy soils, with a good drainage and the soil should warm up quickly. Standard fertilization: 60-80 g/m<sup>2</sup> of a slow release fertilizer.

**Temperature:**

Grow at 15 °C or outdoors. In winter indoors frost free at 3-5 °C or outdoors. Outdoors a fleece cover or mulch cover with straw (20 cm) is needed. In spring the plants start to grow at 12-15 °C and long day. A chilling period for flower initiation is not required, but with a chilling period the flowering is more uniform and full. Kniphofia can also be made to flower 2-4 weeks earlier in the poly-tunnel.

**Fertilization:**

High fertilization levels are required. Fertilize the crop weekly with 100-150 ppm nitrogen (at 3 kg/m<sup>3</sup> slow release fertilizer in substrate), using alternating a complete balanced fertilizer and calcium nitrate fertilizer. Avoid high ammonium

levels. Don't fertilize after mid September. In spring fertilize with 100-150 ppm nitrogen of a potassium balanced fertilizer (N: K<sub>2</sub>O-ratio: 1:1,5). Prevent magnesium deficiency by applying magnesium sulphate (0,05 %) 1-2 times and in case of iron deficiency apply iron-chelate for 1-2 times. Field: If necessary according to analysis, improve the soil with 100-150 g/m<sup>2</sup> of a slow release fertilizer per year, applied in several portions.

Ref. : 8 May 2013  
www.benary.com

